



Ramsey numbers for a disjoint union of some graphs

Halina Bielak

Institute of Mathematics, UMCS, Lublin, Poland

ARTICLE INFO

Keywords:

Complete graph

Forest

G-good graph

Ramsey number

Tree

ABSTRACT

We give the Ramsey number for a disjoint union of some G -good graphs versus a graph G generalizing the results of Stahl [S. Stahl, On the Ramsey number $r(F, K_m)$ where F is a forest, Canad. J. Math. 27 (1975) 585–589] and Baskoro et al. [E.T. Baskoro, Hasmawati, H. Assiyatun, Note. The Ramsey number for disjoint unions of trees, Discrete Math. 306 (2006) 3297–3301].

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

Let G, H, F be simple graphs with at least two vertices. The Ramsey number $R(G, H)$ is the smallest integer n such that every graph F of order n contains a subgraph isomorphic to G or \bar{F} contains a subgraph isomorphic to H , where \bar{F} is the complement of F .

For graphs G, H such that H is a subgraph of G , let us define $G - H$ as the graph obtained from G by deleting the vertices of H and all edges incident to them.

The graph H is G -good if $R(H, G) = (\chi(G) - 1)(|V(H)| - 1) + s(G)$, where $s(G)$ is the chromatic surplus of G , i.e. the minimum cardinality of colour classes over all chromatic colourings of $V(G)$. Let T_i be a tree of order i .

Chvátal [4] proved that $R(T_n, K_m) = (m - 1)(n - 1) + 1$, where T_n is a tree of order n , K_m is a complete graph of order m , with $n, m > 1$. Thus trees are K_m -good graphs.

Let F be a graph, $c(F)$ be the order of the largest component of F and $k_i(F)$ be the number of components of order i in F .

Stahl [7] extended the result of Chvátal for the family of disjoint trees and proved the following theorem.

Theorem 1 (Stahl [7]). *If F is an arbitrary forest then*

$$R(F, K_m) = \max_{1 \leq j \leq c(F)} \left\{ (j - 1)(m - 2) + \sum_{i=j}^{c(F)} ik_i(F) \right\}.$$

Baskoro et al. [1] proved the equivalent form of the formula of Stahl for special cases of forests. Their result is cited below.

Theorem 2 (Baskoro et al. [1]). *Let $n_i \geq n_{i+1}$ for $i = 1, 2, \dots, k - 1$ and $m \geq 2$. If $n_i \geq (n_i - n_{i+1})(m - 1)$ for any i , then*

$$R\left(\bigcup_{i=1}^k T_{n_i}, K_m\right) = R(T_{n_k}, K_m) + \sum_{i=1}^{k-1} n_i.$$

We prove a generalization of both of the above results, where instead of K_m we consider a graph G with $s(G) = 1$, and instead of a family of trees we consider a family of some graphs consisting of G -good components. Further extension of the results to a more general class of graphs with G -good components for $s(G) \geq 1$ is presented in [2].

E-mail address: hbiel@hektor.umcs.lublin.pl.

2. Results

First we prove the following lemma.

Lemma A. Let G be a graph with $\chi(G) = m$ and $s(G) = 1$. If B is a graph with t G -good components, each of them with n vertices, then

$$R(B, G) = (n-1)(m-2) + nt.$$

Proof. The graph $H = K_{nt-1} \cup (m-2)K_{n-1}$ does not contain B . Moreover, $\chi(\bar{H}) \leq m-1$, which means that \bar{H} does not contain G . So we get $R(B, G) \geq (n-1)(m-2) + nt$. The reverse inequality is proved by induction on t . For $t = 1$ the assertion holds since the lemma is reduced to the definition of a G -good graph with $s(G) = 1$.

So assume the assertion of the lemma for all graphs with $t-1$ ($t > 1$) components each of which is a G -good graph on n vertices with $s(G) = 1$.

Let H be a graph on $(n-1)(m-2) + nt$ vertices such that \bar{H} does not contain G . We will prove that H contains B . Let C be a component of B . Since $|V(H)| = (n-1)(m-2) + nt > (n-1)(m-2) + n$ we conclude that H contains C . Note that $|V(H) - V(C)| = (n-1)(m-2) + n(t-1)$. In view of the inductive hypothesis, we get that $H - C$ contains $B - C$. So H contains B . \square

By Lemma A and the idea of the proof for a forest versus a complete graph presented in [7] we get the following more general theorem.

Theorem 3. Let G be a graph with $\chi(G) = m$ and $s(G) = 1$. If B is a graph with G -good components then

$$R(B, G) = \max_{1 \leq j \leq c(B)} \left\{ (j-1)(m-2) + \sum_{i=j}^{c(B)} ik_i(B) \right\}.$$

Proof. Let B_j be the subgraph of B consisting of all the components with at least j vertices, where $1 \leq j \leq c(B)$. Evidently, $B_j - B_{j+1}$ consists of $k_j(B)$ components with exactly j vertices. Moreover, B_{j+1} is the subgraph of B_j . Suppose that the maximum is achieved for $j = j_0$. Set $p_0 = \sum_{i=j_0}^{c(B)} ik_i(B)$. Let $r = (j_0-1)(m-2) + p_0$. Note that the graph $H = K_{p_0-1} \cup (m-2)K_{j_0-1}$ does not contain any subgraph B_{j_0} . So H does not contain the graph B . Moreover, since $\chi(G) = m$ and $\chi(\bar{H}) \leq m-1$, the complement of H does not contain G . Thus $R(B, G) \geq r$. To prove the reverse inequality suppose that the complement of a graph H of order r does not contain G . We shall show that H contains B . Let us assume that $c(B) = n$. Note that $r \geq (n-1)(m-2) + nk_n(B)$. Thus, by Lemma A, we have that H contains B_n . Now we use descending induction to show that H contains B_j for each $j \geq 1$. Let us state the inductive hypothesis: H contains B_{j+1} for some $1 \leq j < n$. Note that B_{j+1} has $\sum_{i=j+1}^n ik_i(B)$ vertices. Thus $H - B_{j+1}$ has $r - \sum_{i=j+1}^n ik_i(B)$ vertices. By the definition of r we get $r \geq (j-1)(m-2) + \sum_{i=j}^n ik_i(B)$. Hence, $r - \sum_{i=j+1}^n ik_i(B) \geq (j-1)(m-2) + jk_j(B) = R(B_j - B_{j+1}, G)$. The above equality follows by Lemma A. So the graph $H - B_{j+1}$ contains $B_j - B_{j+1}$ and therefore H contains B_j . By induction, H contains B_1 . \square

Similarly the result of Baskoro et al. can be extended for graphs G and B such that $s(G) = 1$ and B has G -good components. The theorem below presents a generalization.

Theorem 4. Let G be a graph with $\chi(G) = m$ and $s(G) = 1$. Let $B = \bigcup_{i=1}^k B_{n_i}$, where B_{n_i} is a connected G -good graph of order n_i ($1 \leq i \leq k$). Let for $i = 1, 2, \dots, k-1$, $n_i \geq n_{i+1}$ and $p(n_i) = 1$ if $n_i \geq (n_i - n_{i+1})(m-1)$, and $p(n_i) = 0$ in the opposite case. If $p(n_i) = 0$ for each $j \leq i \leq k-1$ and $p(n_i) = 1$ for $1 \leq i \leq j-1$, where j is an integer $1 \leq j \leq k$, then

$$R(B, G) = (m-1)(n_k-1) + \sum_{i=1}^{k-1} n_i p(n_i) + (m-1) \sum_{i=1}^{k-1} (n_i - n_{i+1})(1 - p(n_i)) + 1.$$

Proof. Let $t = (m-1)(n_k-1) + \sum_{i=1}^{k-1} n_i p(n_i) + (m-1) \sum_{i=1}^{k-1} (n_i - n_{i+1})(1 - p(n_i)) + 1$.

First we prove the inequality $R(B, G) \leq t$ by induction on k . For $k = 1$ the result is immediate.

Let us assume the inductive hypothesis for all graphs consisting of s components (each being a G -good graph), where $1 \leq s < k$:

$$R\left(\bigcup_{i=1}^s B_{n_i}, G\right) = (m-1)(n_s-1) + \sum_{i=1}^{s-1} n_i p(n_i) + (m-1) \sum_{i=1}^{s-1} (n_i - n_{i+1})(1 - p(n_i)) + 1.$$

Let us take an arbitrary graph H of order t . Suppose that \bar{H} does not contain G as a subgraph. Note that

$$t = (m-1)(n_{k-1}-1) + 1 + \sum_{i=1}^{k-2} n_i p(n_i) + (m-1) \sum_{i=1}^{k-2} (n_i - n_{i+1})(1 - p(n_i)) + t_0,$$

where

$$t_0 = -(m-1)(n_{k-1} - n_k) + n_{k-1}p(n_{k-1}) + (m-1)(n_{k-1} - n_k)(1 - p(n_{k-1})).$$

Note that $t_0 \geq 0$. So, by the inductive hypothesis, H contains $F = \bigcup_{i=1}^{k-1} B_{n_i}$.

Note that $t - \sum_{i=1}^{k-1} n_i \geq (m-1)(n_k - 1) + 1$. So the graph $H - F$ contains B_{n_k} . Hence, H contains $B = \bigcup_{i=1}^k B_{n_i}$.

To prove the inequality $R(B, G) \geq t$ let us define a graph H of order $t - 1$ such that H does not contain B as a subgraph and \bar{H} does not contain G as a subgraph.

We first note that $t - 1 = (m-2)x + y$, where $x = n_k - 1 + \sum_{i=1}^{k-1} (n_i - n_{i+1})(1 - p(n_i))$ and $y = \sum_{i=1}^k n_i - 1 - \sum_{i=1}^{k-1} n_{i+1}(1 - p(n_i))$. Moreover, by the assumption of the theorem, we have $x = n_j - 1$ and $y = \sum_{i=1}^j n_i - 1$.

Thus, let us consider the graph $H = (m-2)K_x \cup K_y$. Note that $B_{n_i} \not\subset K_x$ for $i \leq j$.

If $B \subset H$, then $\bigcup_{i=1}^j B_{n_i} \subset K_y$. But this is impossible because $\sum_{i=1}^j n_i > y$. Finally, $G \not\subset \bar{H}$ (since $\chi(G) - 1 = \chi(\bar{H})$). \square

3. Remarks

Finally, let us consider some examples of the graphs consisting of G -good components, where $s(G) = 1$. Let $B = T_7 \cup T_5 \cup T_2$ and $F = T_5 \cup T_2 \cup T_2$, where T_n is a tree of order n . Each component of the graphs B and F is K_3 -good. Note that $p(n_1) = 1$ and $p(n_2) = 0$ for the graph B . Thus the condition of Bascoro et al. does not hold for B . The assumption of Theorem 4 holds and we can apply the respective formula for counting $R(B, K_3) = 16$ (we can use Theorem 3, as well). Then note that $p(n_1) = 0$ and $p(n_2) = 1$ for the graph F . Thus the assumption of Theorem 4 does not hold for F and K_3 . Evidently, the condition of Bascoro et al. does not hold as well. By Theorem 3, we have $R(B, K_3) = 10$. Note that the right hand side of the formula in Theorem 4 is equal to 11 for this pair of graphs. Similarly, the components of the graphs $B = C_{12} \cup C_5 \cup C_5$ and $F = C_{23} \cup C_{12} \cup C_5 \cup C_5$ are C_5 -good, where C_n is a cycle of order n . So, by Theorem 3, we get $R(B, C_5) = 26$ and $R(F, C_5) = 48$ and the right hand of the formula in Theorem 4 is 28 and 50, respectively. In fact, the graphs B and F do not satisfy the assumption of Theorem 4. Evidently, the assumption holds for all graphs B with G -good components, where $\chi(G) = 2$ and $s(G) = 1$.

Problem. We can formulate the following problem. Characterize the sequences of orders of the components of a graph B for which the formula of Theorem 4 holds. Evidently, if each component of a graph B is G -good and $s(G) = 1$, then the right hand of the formula in Theorem 4 is the upper bound for $R(B, G)$ (see the first part of the proof of Theorem 4).

One can find some other examples of G -good graphs with $s(G) = 1$ in [3,5,6].

An attractive collection of applications of various branches of Ramsey theory in algebra, geometry and point-set topology is presented in B. Bollobás, Graph Theory, Springer GTM63, 1979.

Acknowledgments

The author is grateful to anonymous referees for their helpful suggestions on the presentation of the work.

References

- [1] E.T. Baskoro, Hasmawati, H. Assiyatun, Note. The Ramsey number for disjoint unions of trees, Discrete Math. 306 (2006) 3297–3301.
- [2] H. Bielak, Ramsey numbers for disjoint union of good graphs (submitted for publication).
- [3] Y. Chen, Y. Zhang, K. Zhang, The Ramsey numbers of trees versus W_6 or W_7 , European J. Comb. 27 (2006) 558–564.
- [4] V. Chvátal, Tree-complete graph Ramsey number, J. Graph Theory 1 (1977) 93.
- [5] A.N.M. Salman, H.J. Broersma, The Ramsey numbers of paths versus kipases, Electron. Notes Discrete Math. 17 (2004) 251–255.
- [6] A.N.M. Salman, H.J. Broersma, Path-fan Ramsey numbers, Discrete Appl. Math. 154 (2006) 1429–1436.
- [7] S. Stahl, On the Ramsey number $r(F, K_m)$ where F is a forest, Canad. J. Math. 27 (1975) 585–589.